






Report

Skin cancer mortality in Spain: adjusted mortality rates by province and related risk factors

Antonio Alcalá Ramírez del Puerto¹, MD, Juan Carlos Hernández-Rodríguez¹, MD, 
Mercedes Sendín-Martín¹, MD,  Juan Ortiz-Alvarez¹, MD, 
Julián Conejo-Mir Sánchez^{1,2},  and José Juan Pereyra-Rodríguez^{1,2}, MD, PhD 

¹Dermatology Department, Hospital Universitario Virgen del Rocío, Sevilla, Spain and ²School of Medicine, Medicine Department, Sevilla University, Sevilla, Spain

Correspondence

José-Juan Pereyra-Rodríguez, PhD
H.U. Virgen del Rocío
Avda. Manuel Siurot s/n
41013 Sevilla
Spain
E-mail: jpereyra@us.es

Conflict of interest: None.

Funding source: None.

doi: 10.1111/ijd.16618

Abstract

Background Ultraviolet radiation is the main environmental risk factor responsible for the development of skin cancer. Other occupational, socioeconomic, and environmental factors appear to be related to the risk of skin cancer. Furthermore, the factors appear to differ for melanoma and non-melanoma skin cancer (NMSC). The purpose of this study is to analyze mortality rates of skin cancer in the different provinces of Spain and to determine the influence of socioeconomic conditions and other environmental and demographic factors in rates.

Methods Deaths from melanoma and NMSC in the period 2000–2019 were obtained as well as socioeconomic and environmental variables. Annual standardized mortality rates (SMR) were calculated for all Spanish provinces. The Pearson correlation coefficient was calculated.

Results The SMR of melanoma was 2.10/100,000 inhabitants, while that of NMSC was 1.28/100,000. At the provincial level, a great variability is confirmed. Gross domestic product showed a positive correlation with melanoma mortality but a negative correlation with NMSC. Other environmental and socioeconomic variables also showed correlation, as a positive correlation between tobacco sales and melanoma and between agricultural development and the NMSC.

Conclusions There are still important differences between each province that must be taken into account when planning health care and resource distribution. This ecological and province-wise study helps to elucidate the relationship between social and ambient exposure determinants and skin cancer mortality in Spain.

Introduction

Skin cancer and its three main subtypes, basal cell carcinoma (BCC), squamous cell carcinoma (SCC), and malignant melanoma (MM), represent the most common form of cancer among Caucasians worldwide.¹ Ultraviolet radiation is the main environmental risk factor responsible for the development of skin cancer, with the most common subtype of MM, superficial spreading melanoma, being related to intermittent excessive sun exposure and sunburn in childhood, while SCC has been associated with chronic exposure to ultraviolet radiation.^{2–5} However, increased sun exposure does not always imply a greater risk of developing skin cancer, as numerous studies and meta-analyses show how chronic occupational sun exposure seems to be inversely associated with the development of MM, playing a protective role.^{6,7}

Melanoma and non-melanoma skin cancer (NMSC), corresponding mostly to SCC and BCC, and to a lesser extent other

types such as Merkel or cutaneous sarcomas, have maintained a growing incidence since the end of the last century, reaching an alarming annual increase of up to 7% in the case of melanoma, although mortality rates have remained stable or even decreasing, this decrease in mortality being possible because of the introduction of targeted immunotherapy.^{8,9} The increased incidence is probably related to factors that directly influence sun exposure such as the increase in outdoor recreational activities, changes in lifestyle and behavior toward the sun, increase in coastal tourism, or the persistence of the tanning culture, although other unrelated factors may come into play directly with sun exposure such as genetics, population aging, and the increase in the immunosuppressed population.^{10,11}

Due to the cultural influence on sun exposure, numerous studies have been conducted globally on the influence of socioeconomic factors on the incidence and mortality of skin cancer, especially in MM, and most of them reaffirm how there is greater incidence but lower mortality in populations with greater

economic wealth, possibly because of the greater culture they have for recreational sun exposure habits and earlier dermatological care.¹² However, no similar study has been carried out at the national level in Spain.

The purpose of this study is to analyze how mortality rates behave in the different types of skin cancer in the different provinces of Spain in the period 2000 to 2019, and, parting from the same database, to determine the influence of socioeconomic conditions and other environmental and demographic factors in the mortality rate of MM and NMSC at the national level.

Materials and methods

We have carried out an ecological study that included deaths from melanoma and non-melanoma skin cancer in the different Spanish provinces in the period 2000–2019, obtained from microdata files provided by the National Institute of Statistics¹³ (CIE10 codes C43 [MM] and C44 [NMSC]). Population data for each province at mid-year for the same period of years.¹³ Annual standardized mortality rates (SMR) were calculated using the direct method for all Spanish provinces, using the 2013 European standard population as a reference and using the Epidat 4.2 program. The results obtained were expressed as deaths per 100,000 and with a 95% CI. Subsequently, the different provincial variables with which the SMR are compared were obtained through the National Institute of Statistics,¹³ the State Meteorological Agency,¹⁴ the National Catalog of

Hospitals,¹⁵ the Tax Agency,¹⁶ and public data from the Tobacco Market.¹⁷ The Pearson correlation coefficient was calculated with the SPSS V25 program between the SMR of each province and the variables collected annually at the provincial level (number of hospital beds; level of agricultural development [defined as the percentage of workers employed in the agriculture sector by province]; gross domestic product; total tobacco sold by each province; geographical and meteorological variables such as altitude, longitude, and latitude; annual mean irradiation and insolation; annual mean, maximum, and minimum temperature; annual absolute minimum and maximum temperature; annual highest normal minimum temperature; annual lowest maximum normal temperature; and number of rainy and foggy days in the year). A significant correlation was considered if it reached an absolute r greater than 0.3. To create the map with the color gradient by province, for melanoma, and for NMSC, Microsoft Excel 2016 program was used.

Results

At the national level, the SMR of MM was 2.10/100,000 inhabitants, while that of NMSC was 1.28/100,000. At the provincial level, a great variability is confirmed, with the highest SMR of MM in Albacete (2.57) and Álava (2.48) and of NMSC in Ceuta (2.34) and Soria (2.15). The SMR of each province is shown in Figures 1 and 2 (Tables 1 and 2).




Figure 1 Standardized annual melanoma mortality rates by province, 2000–2019




Figure 2 Standardized annual SCC mortality rates by province, 2000–2019

Table 1 Age-specific and standardized melanoma mortality rate per 100,000 inhabitants (European standard population) in Spain

Age (years)	Both genders				Males				Females			
	2000–2004	2005–2009	2010–2014	2015–2019	2000–2004	2005–2009	2010–2014	2015–2019	2000–2004	2005–2009	2010–2014	2015–2019
0–4	0	0	0.02	0	0	0	0.02	0	0	0	0.02	0
5–9	0.01	0	0	0.01	0	0	0	0.02	0.02	0	0	0
10–14	0.02	0.01	0.03	0	0.02	0.02	0.02	0	0.02	0	0.04	0
15–19	0.04	0.03	0.03	0.01	0.02	0.03	0.05	0	0.07	0.04	0	0.02
20–24	0.17	0.11	0.06	0.07	0.2	0.1	0.05	0	0.14	0.13	0.08	0.14
25–29	0.26	0.21	0.16	0.32	0.28	0.23	0.23	0.31	0.23	0.19	0.09	0.33
30–34	0.58	0.42	0.42	0.39	0.63	0.48	0.46	0.43	0.53	0.36	0.37	0.35
35–39	0.84	0.69	0.66	0.68	0.85	0.77	0.68	0.74	0.84	0.61	0.63	0.63
40–44	1.02	1.11	0.87	0.86	1.15	1.21	0.93	0.95	0.89	0.99	0.81	0.77
45–49	1.56	1.37	1.39	1.24	1.7	1.45	1.53	1.48	1.43	1.28	1.25	0.99
50–54	1.97	1.87	1.74	1.71	2.52	2.15	2.06	1.88	1.44	1.6	1.41	1.54
55–59	2.42	2.48	2.4	2.16	3.07	2.98	3.03	2.53	1.81	2	1.8	1.79
60–64	3.14	3.16	3.17	2.89	4.32	4.05	4.06	3.75	2.05	2.32	2.32	2.07
65–69	4.12	4.23	4.32	3.91	4.94	5.41	5.64	5.1	3.4	3.16	3.12	2.82
70–74	5.01	5.55	5.63	5.71	6.61	7.18	8.08	7.66	3.71	4.21	3.56	4.02
75–79	6.04	6.79	7.61	7.22	7.61	9.17	10.32	10.23	4.92	5.03	5.54	4.86
80–84	7.51	8.84	9.18	10.87	10.3	11.64	13.36	14.65	5.89	7.09	6.44	8.31
>85	12.68	12.19	13.29	14.31	14.76	16.12	17.47	18.63	11.78	10.45	11.31	12.13
SMR	2.01	2.06	2.1	2.1	2.5	2.6	2.75	2.7	1.64	1.65	1.6	1.63

SMR, standardized mortality rates.

Table 2 Age-specific and standardized NMSC mortality rate per 100,000 inhabitants (European standard population) in Spain

Age (years)	Both genders				Males				Females			
	2000–2004	2005–2009	2010–2014	2015–2019	2000–2004	2005–2009	2010–2014	2015–2019	2000–2004	2005–2009	2010–2014	2015–2019
0–4	0	0	0	0	0	0	0	0	0	0	0	0
5–9	0	0.01	0	0	0	0	0	0	0	0.02	0	0
10–14	0	0	0	0	0	0	0	0	0	0	0	0
15–19	0.01	0.01	0	0	0.02	0.02	0	0	0	0	0	0
20–24	0.03	0.01	0.01	0.02	0.02	0	0.02	0	0.04	0.01	0	0.04
25–29	0.04	0.02	0.05	0.02	0.07	0.01	0.04	0	0.01	0.02	0.05	0.03
30–34	0.05	0.03	0.04	0.03	0.06	0.05	0.03	0.05	0.04	0.01	0.05	0.01
35–39	0.08	0.06	0.05	0.04	0.11	0.08	0.07	0.03	0.05	0.04	0.03	0.04
40–44	0.1	0.11	0.15	0.05	0.13	0.14	0.18	0.05	0.08	0.08	0.12	0.04
45–49	0.22	0.1	0.2	0.11	0.3	0.13	0.22	0.14	0.15	0.07	0.19	0.09
50–54	0.3	0.25	0.3	0.24	0.49	0.38	0.49	0.33	0.11	0.13	0.11	0.16
55–59	0.49	0.6	0.55	0.37	0.77	0.97	0.78	0.58	0.22	0.25	0.33	0.16
60–64	0.72	0.72	0.81	0.67	1.07	1.1	1.12	1.1	0.4	0.36	0.52	0.28
65–69	1.15	1.37	1.26	1.13	1.96	2.28	1.97	1.82	0.44	0.56	0.62	0.51
70–74	2.01	1.8	2.06	1.88	3.13	2.83	3.4	3.32	1.1	0.96	0.92	0.65
75–79	3.39	3.66	3.91	3.54	5.3	6.06	6.6	6.1	2.02	1.89	1.86	1.54
80–84	7.94	7.52	7.42	7.44	11.66	12.32	12	12.47	5.78	4.53	4.41	4.02
>85	31.87	26.16	26.55	26.87	42.59	34.53	37.94	38.19	27.26	22.45	21.15	21.18
SMR	1.43	1.28	1.32	1.25	2.03	1.87	1.98	1.92	1.06	0.89	0.88	0.81

SMR, standardized mortality rates.

A positive correlation was obtained between the MM SMR and the provincial gross domestic product (GDP) ($r = 0.47$), and the sale of tobacco in each province ($r = 0.33$) and negative with geographic longitude ($r = -0.47$; Table 3).

Regarding NMSC, a positive correlation was obtained between the NMSC SMR and the level of provincial agricultural development ($r = 0.35$), mean annual temperature ($r = 0.41$), maximum mean annual temperature ($r = 0.41$), annual mean minimum temperature ($r = 0.36$), absolute minimum temperature ($r = 0.31$), highest minimum temperature of the year ($r = 0.40$), and lowest maximum temperature of the year ($r = 0.38$). A negative correlation was obtained between NMSC mortality and the number of provincial hospital beds ($r = -0.40$), provincial GDP ($r = -0.50$), and latitude ($r = -0.40$) (Table 3).

Discussion

In Europe, melanoma incidence rates follow a north-south gradient, with the highest rates in northern European countries. This distribution is similar in the rest of the world, with a higher incidence rate as we move away from the equator. In Spain, a crude global incidence of 8.82 cases per 100,000 people/year has been estimated, with a crude global mortality of 2.17 cases per 100,000 people/year and a low standardized incidence rate compared to other countries in Europe, the USA, Australia, and New Zealand.¹⁸

Regarding SCC, in a recent 2016 meta-analysis,¹⁹ it was estimated that the crude incidence rate in Spain was 38.16 per

100,000 person-years, our country being in the low range of incidence along with countries such as Germany or Slovakia. The mortality of SCC, although low, represents the highest percentage of it for NMSC.

The melanoma mortality at the national level obtained in our analysis (2.10/100,000 inhabitants) is very similar to that obtained in recent studies (2.17/100,000 inhabitants¹⁹ and 2.13/100,000 inhabitants,⁹ respectively). Despite the fact that the mortality gradient due to MM in Europe is well-known, with higher mortality in the southern and eastern countries,²⁰ our results have shown that Spain continues to be an exception, possibly explained by the protective role that plays in chronic sun exposure, the higher skin phototype, and universal and free access to the health system.²¹ To our knowledge, this is the first study to analyze mortality at the provincial level. It highlights a higher mortality in the eastern provinces of the country, a fact confirmed by the statistically significant positive correlation between SMR and geographical longitude ($r = -0.47$, $P = 0.001$). This phenomenon could be because of multiple factors, such as the greater beach culture of these provinces and, consequently, the greater intermittent sun exposure that predisposes to the development of MM, as well as the greater number of foreigners with a lower skin phototype. Paradoxically, contrary to what might be expected, no correlation was observed with the number of hospital beds and the mean irradiation, and the mean insolation.

However, the standardized NMSC mortality rate obtained (1.28 per 100,000 inhabitants) is practically identical to that

Table 3 Pearson correlation coefficient (bilateral significance) between adjusted mortality rates for melanoma and non-melanoma skin cancer and multiple socioeconomic and environmental variables by province during the period 2000–2019

	NMSC 2000–2019	Melanoma 2000–2019
Hospital beds	−0.40**	0.16
Agricultural development	0.35*	−0.26
GDP	−0.50**	0.47**
Tobacco	−0.26	0.33*
Altitude	−0.14	0.06
Longitude	0.04	−0.47**
Latitude	−0.40**	0.22
Irradiation	0.40	0.16
Insolation	0.266	−0.113
TM_MAX	0.41**	−0.23
TM_MES	0.41**	−0.20
TM_MIN	0.36*	−0.14
TA_MAX	0.24	−0.28*
TA_MIN	0.31*	−0.17
TS_MIN	0.40**	−0.12
TI_MAX	0.38**	−0.22
N_LLU	−0.29*	0.04
N_FOG	−0.28*	−0.05

NMSC, non-melanoma skin cancer; GDP, gross domestic product; TM_MAX, average annual maximum temperature (°C); TM_MES, average annual temperature (°C); TM_MIN, average annual minimum temperature (°C); TA_MAX, annual absolute maximum temperature (°C); TA_MIN, annual absolute minimum temperature (°C); TS_MIN, annual highest normal minimum temperature (°C); TI_MAX, annual lowest maximum normal temperature (°C); N_LLU, number of rainy days in the year; N_FOG, number of foggy days in the year.

* < 0.05 .

** < 0.01 .

described by Sendin et al.²² which was 1.27 per 100,000 inhabitants in 2018 and slightly higher than the one reported in a 2016 meta-analysis¹⁹ (1.10 per 100,000 person-years). The mortality rate of NMSC is low; it is to a greater extent caused by the high-risk variant of SCC and is similar to that of other European countries.^{23–25} At the provincial level, higher mortality from NMSC is found in provinces with greater agricultural development ($r = 0.35$, $P = 0.024$), with higher average temperature, and in those with lower latitude ($r = 0.4$, $P = 0.004$). Provincial agricultural development and average temperature can be seen as indirect markers of chronic exposure to the sun, which is well-known to be an important risk factor for the development of SCC.⁴ The statistically significant negative correlation obtained with the meteorological variables that are inversely related to chronic solar radiation, such as the number of days of fog or rain, reinforces the theory of association between chronic sun exposure and SCC.

It is worth noting the correlation that has been obtained with provincial GDP (positive with SMR MM, $r = 0.47$ and negative

with SMR NMSC, $r = -0.50$). If we assume that provincial GDP is an indirect measure of provincial per capita income and economic wealth, these results contradict what is described in the literature. At a global level, numerous articles have been written on the influence of socioeconomic factors on the incidence and mortality of skin cancer, especially in MM. In the USA, Singh et al. in 2011²⁶ observed a higher incidence rate of MM in regions with less poverty, higher education, higher income, and lower unemployment. However, this higher incidence does not correspond to higher mortality, since Sitenga et al. in 2018²⁷ corroborated how the populations with the lowest income in the US percentile have lower survival and greater probability of dying from MM in all its stages, probably because of the delay in consulting a specialist and the difficulty of access to medical care in privatized health systems. At a European level, similar studies have been carried out in Denmark,²⁸ the Netherlands²⁴ and other northern countries²⁹ that reaffirm the higher incidence in populations with greater economic wealth, possibly because of intermittent recreational exposure to the sun of typical hobbies of higher economic status, coastal vacations in childhood, and a greater culture of tanning, activities that have been shown to increase the risk of MM.^{6,12,30} Internationally, mortality and affluence show a similar relationship with patients with lower socioeconomic status (SES) having lower incidence^{31–33} but being diagnosed at a later stage. In a 2015 meta-analysis where 32 studies were analyzed,³⁴ it was observed that although higher SES populations generally have a higher incidence of melanoma, lower SES populations have higher mortality from melanoma. However, in Spain at an ecological level, we observe that there is a higher mortality from MM in the provinces with higher income. This may be due to several factors, the main being the lower incidence of nodular melanoma that exists in Spain with respect to other nearby countries.³⁵ This subtype has been associated in several studies with a lower SES.^{31,36} Other factors may also have contributed to explaining this difference, such as universal and free access to the healthcare system and the ecological nature of our study. Further analytical studies at the national level are needed to clarify this issue.

The relationship between NMSC and socioeconomic status has less been studied. Steding et al. in 2010³⁷ observed a higher incidence of SCC in the rural population with lower incomes, as well as higher mortality.³⁸ Similar results have been obtained in our analysis, obtaining a positive correlation with the agricultural development of each province. This may be explained by the fact that living in the countryside leads to an accumulation of smaller doses of UVR day by day, ultimately to a higher cumulated dose of UVR, and consequently to a higher risk of SCC. Likewise, this more chronic exposure would explain the lack of correlation with deaths from melanoma. The same can be said of the existing negative correlation with the number of hospital beds, since this is still an indirect measure of the economic development of each province.

The association between smoking and skin cancer is controversial. Our study obtained a positive correlation between melanoma mortality and provincial tobacco sales ($r = 0.33$), while no significant correlation was obtained with NMSC mortality. A recent meta-analysis³⁹ showed that smoking could moderately increase the risk of SCC but slightly reduce the risk of MM, contrary to our results. It may be explained as smoking encompasses many carcinogenic compounds, attenuates immune responses, and decreases the cutaneous blood flow, which increases the risk of SCC among smokers.^{40,41} Smoking also hinders the growth of melanoma cells by downregulating the gene expression of the Notch pathway, which controls cell differentiation and produces the accumulation of nicotine in tissues containing melanin, thus suppressing inflammatory responses to UV radiation.^{42,43} However, in agreement with our results, Pozniak et al.⁴⁴ observed a higher mortality from melanoma in heavy smokers. In this study, smoking had an adverse effect on melanoma-specific outcome by reducing the protective value of immune infiltration, with a greater detrimental effect of smoking on survival with the strength of the tumor immune signal of the patient. Smoking can also be a detriment to healing from skin cancer surgery. In fact, numerous scientific studies have shown that smoking cigarettes can significantly reduce the blood flow necessary for wounds to heal properly.⁴⁵ Finally, in another study carried out in Spain, using a propensity score matching analysis, the authors found no impact of smoking on survival of melanoma patients and that it should not influence treatment decisions.⁴⁶ With all these data, and given that our data show an association between smoking and mortality, we hypothesize that the smoking patients included in our study could present more aggressive melanomas.

The fundamental limitation of this study is that of analyses at the ecological level, where we can only establish correlations between variables, while the individual distribution of exposure and health remains unknown (ecological fallacy), which leads to a possible distortion of the association between exposure and outcome. However, they are an excellent way to establish hypotheses that can be confirmed later at the individual level.

Conclusion

The burden of skin cancer presents an urgent need for increased early detection and public education. Although skin cancer mortality in Spain is low compared to other nearby countries and has decreased in recent years, there are still important differences between each province that must be taken into account when planning health care and resource distribution. This ecological and province-wise study helps to elucidate the relationship between social and ambient exposure determinants and skin cancer mortality in Spain, with a higher mortality of melanoma in provinces of higher income and higher mortality of NMSC in the poorer and more agricultural-oriented provinces. To our knowledge, this is the first Spanish study that takes into

account every province and may provide the foundation for more analytical studies to help clarify the complex relationship between skin cancer mortality and socioeconomic and environmental factors, leading to more effective targeted interventions and education on sun habits at the individual, community, province, and national level.

References

- 1 Tejera-Vaquerizo A, Descalzo-Gallego MA, Otero-Rivas MM, Posada-García C., Rodríguez-Pazos L, Pastushenko I, et al. Incidencia y mortalidad del cáncer cutáneo en España: revisión sistemática y metaanálisis. *Actas Dermosifiliogr*. 2016;**107**:318–28.
- 2 Brunssen A, Waldmann A, Eisemann N, Katalinic A. Impact of skin cancer screening and secondary prevention campaigns on skin cancer incidence and mortality: a systematic review. *J Am Acad Dermatol*. 2017;**76**:129–139.e10.
- 3 Armstrong BK, Kricke A. How much melanoma is caused by sun exposure? *Melanoma Res*. 1993;**3**:395–401.
- 4 Armstrong BK, Cust AE. Sun exposure and skin cancer, and the puzzle of cutaneous melanoma: a perspective on Fears et al. Mathematical models of age and ultraviolet effects on the incidence of skin cancer among whites in the United States. *American Journal of Epidemiology* 1977. *Cancer Epidemiol*. 2017;**48**:147–56.
- 5 Carr S, R. F. of M, Smith C, Wernberg J. Epidemiology and risk factors of melanoma. *Surg Clin North Am*. 2020;**100**:1–12.
- 6 Gandini S, Sera F, Cattaruzza MS, Pasquini P, Picconi O, Boyle P, et al. Meta-analysis of risk factors for cutaneous melanoma: II. Sun exposure. *Eur J Cancer*. 2005;**41**:45–60.
- 7 Zanetti R, Rosso S, Martinez C, Navarro C, Schraub S, Sancho-Garnier H, et al. The multicentre south European study 'Helios'. I: skin characteristics and sunburns in basal cell and squamous cell carcinomas of the skin. *Br J Cancer*. 1996;**73**:1440–6.
- 8 Martín García E, Arias-Santiago S, Serrano-Ortega S, Buendía-Eisman A. Changes in the incidence of skin and lip cancer between 1978 and 2007. *Actas Dermosifiliogr*. 2017;**108**:335–45.
- 9 Durán-Romero AJ, Sendín-Martin M, Conejo-Mir J, Pereyra-Rodríguez JJ. Cutaneous malignant melanoma mortality in Spain from 1979 to 2018. Trends and new perspectives in the immunotherapy era. *J Eur Acad Dermatol Venereol*. 2021;**35**:884–91.
- 10 Leiter U, Eigentler T, Garbe C. Epidemiology of skin cancer. *Adv Exp Med Biol*. 2014;**810**:120–40.
- 11 Kojo K, Jansen CT, Nybom P, Huurto L, Laihia J, Ilus T, et al. Population exposure to ultraviolet radiation in Finland 1920–1995: exposure trends and a time-series analysis of exposure and cutaneous melanoma incidence. *Environ Res*. 2006;**101**:123–31.
- 12 Cortez JL, Vasquez J, Wei ML. The impact of demographics, socioeconomic, and health care access on melanoma outcomes. *J Am Acad Dermatol*. 2020;**84**:1677–83. <https://doi.org/10.1016/j.jaad.2020.07.125>
- 13 Instituto Nacional de Estadística. INEbase. <https://www.ine.es/dyngs/INEbase/listaoperaciones.htm>
- 14 Agencia Estatal de Meteorología. AEMET OpenData. https://www.aemet.es/es/datos_abiertos/AEMET_OpenData
- 15 Ministerio de Sanidad. Catálogo Nacional de Hospitales. <https://www.sanidad.gob.es/ciudadanos/prestaciones/centrosServiciosSNS/hospitales/home.htm>

- 16 Agencia Tributaria. Estadísticas. <https://sede.agenciatributaria.gob.es/Sede/estadisticas.html>
- 17 Ministerio de Hacienda. Información estadística sobre el mercado de tabacos. <https://www.hacienda.gob.es/es-ES/AreasTematicas/CMTabacos/Paginas/EstadisticassobreelMercadodeTabacos.aspx>
- 18 Tejera-Vaquerizo A, Boada A, Nagore E. Why is the cutaneous melanoma mortality rate not falling? *Actas Dermosifiliogr*. 2020;**111**:450–2.
- 19 Tejera-Vaquerizo A, Descalzo-Gallego MA, Otero-Rivas MM, Posada-García C, Rodríguez-Pazos L, Pastushenko I, et al. Cancer incidence and mortality in Spain: a systematic review and meta-analysis. *Actas Dermosifiliogr*. 2016;**107**:318–28.
- 20 Marcoval J, Moreno A, Torras A, Baumann E, Graells J, Gallego MI. Changes in incidence of malignant melanoma in the last 19 years in a tertiary hospital on the mediterranean coast. *Actas Dermo-Sifiliográficas (English Ed)*. 2008;**99**:464–8.
- 21 Forsea AM, Del Marmol V, Stratigos A, Geller AC. Melanoma prognosis in Europe: far from equal. *Br J Dermatol*. 2014;**171**:179–82.
- 22 Sendín-Martin M, Hernández-Rodríguez JC, Durán-Romero AJ, Ortiz-Álvarez J, Conejo-Mir J, Pereyra-Rodríguez JJ. Non-melanoma skin cancer mortality in Spain: a predictive model up to 2044. *J Clin Med*. 2021;**10**(24):5750.
- 23 Wehner MR, Cidre Serrano W, Nosrati A, Schoen PM, Chren MM, Boscardin J, et al. All-cause mortality in patients with basal and squamous cell carcinoma: a systematic review and meta-analysis. *J Am Acad Dermatol*. 2018;**78**:663–672.e3.
- 24 van der Aa MA, de Vries E, Hoekstra HJ, Coebergh JWW, Siesling S. Sociodemographic factors and incidence of melanoma in the Netherlands, 1994–2005. *Eur J Cancer*. 2011;**47**:1056–60.
- 25 Talbäck M, Stenbeck M, Rosén M. Up-to-date long-term survival of cancer patients: an evaluation of period analysis on Swedish Cancer Registry data. *Eur J Cancer*. 2004;**40**:1361–72.
- 26 Singh SD, Ajani UA, Johnson CJ, Roland KB, Eide M, Jemal A, et al. Association of cutaneous melanoma incidence with area-based socioeconomic indicators-United States, 2004–2006. *J Am Acad Dermatol*. 2011;**65**:S58–68.
- 27 Sitenga JL, Aird G, Ahmed A, Walters R, Silberstein PT. Socioeconomic status and survival for patients with melanoma in the United States: an NCDB analysis. *Int J Dermatol*. 2018;**57**:1149–56.
- 28 Birch-Johansen F, Hvilsom G, Kjaer T, Storm H. Social inequality and incidence of and survival from malignant melanoma in a population-based study in Denmark, 1994–2003. *Eur J Cancer*. 2008;**44**:2043–9.
- 29 Idorn LW, Wulf HC. Socioeconomic status and cutaneous malignant melanoma in Northern Europe. *Br J Dermatol*. 2014;**170**:787–93.
- 30 Holman CD, Armstrong BK, Heenan PJ. Relationship of cutaneous malignant melanoma to individual sunlight-exposure habits. *J Natl Cancer Inst*. 1986;**76**:403–14.
- 31 Quintella Mendes GL, Koifman S. Socioeconomic status as a predictor of melanoma survival in a series of 1083 cases from Brazil: just a marker of health services accessibility? *Melanoma Res*. 2013;**23**:199–205.
- 32 Eriksson H, Lyth J, Månsson-Brahme E, Frohm-Nilsson M, Ingvar C, Lindholm C, et al. Low level of education is associated with later stage at diagnosis and reduced survival in cutaneous malignant melanoma: a nationwide population-based study in Sweden. *Eur J Cancer*. 2013;**49**:2705–16.
- 33 Downing A, Newton-Bishop JA, Forman D. Recent trends in cutaneous malignant melanoma in the Yorkshire region of England; incidence, mortality and survival in relation to stage of disease, 1993–2003. *Br J Cancer*. 2006;**95**:91–5.
- 34 Jiang AJ, Rambhatla PV, Eide MJ. Socioeconomic and lifestyle factors and melanoma: a systematic review. *Br J Dermatol*. 2015;**172**:885–915.
- 35 Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 2018;**68**:394–424.
- 36 Pollitt RA, Clarke CA, Swetter SM, Peng DH, Zadnick J, Cockburn M. The expanding melanoma burden in California hispanics: importance of socioeconomic distribution, histologic subtype, and anatomic location. *Cancer*. 2011;**117**:152–61.
- 37 Steding-Jessen M, Birch-Johansen F, Jensen A, Schüz J, Kjaer SK, Dalton SO. Socioeconomic status and non-melanoma skin cancer: a nationwide cohort study of incidence and survival in Denmark. *Cancer Epidemiol*. 2010;**34**:689–95.
- 38 Hannuksela-Svahn A, Pukkala E, Karvonen J. Basal cell skin carcinoma and other nonmelanoma skin cancers in Finland from 1956 through 1995. *Arch Dermatol*. 1999;**135**:781–6.
- 39 Arafa A, Mostafa A, Navarini AA, Dong J-Y. The association between smoking and risk of skin cancer: a meta-analysis of cohort studies. *Cancer Causes Control*. 2020;**31**:787–94.
- 40 D'Errico M, Calcagnile A, Iavarone I, Sera F, Baliva G, Chinni LM, et al. Factors that influence the DNA repair capacity of normal and skin cancer-affected individuals. *Cancer Epidemiol Biomarkers Prev Oncol*. 1999;**8**:553–9.
- 41 De Hertog SA, Wensveen CA, Bastiaens MT, Kielich CJ, Berkhout MJ, Westendorp RG, et al. Relation between smoking and skin cancer. *J Clin Oncol*. 2001;**19**:231–8.
- 42 Sopori M. Effects of cigarette smoke on the immune system. *Nat Rev Immunol*. 2002;**2**:372–7. <https://doi.org/10.1038/nri803>
- 43 Yerger VB, Malone RE. Melanin and nicotine: a review of the literature. *Nicotine Tob Res*. 2006;**8**:487–98.
- 44 Poźniak J, Nsengimana J, Laye JP, O'Shea SJ, Diaz JMS, Droop AP, et al. Genetic and environmental determinants of immune response to cutaneous melanoma. *Cancer Res*. 2019;**79**:2684–96.
- 45 Wang G, Pan C, Cao K, Zhang J, Geng H, Wu K, et al. Impacts of cigarette smoking on the tumor immune microenvironment in esophageal squamous cell carcinoma. *J Cancer*. 2022;**13**:413–25.
- 46 Tejera-Vaquerizo A, Descalzo-Gallego MA, Traves V, Requena C, Bolumar I, Pla A, et al. No association between smoking and sentinel lymph node metastasis and survival in cutaneous melanoma. *J Eur Acad Dermatol Venereol*. 2019;**33**:2283–90.